

SCREENING INSTRUMENT FOR SUBSTANCE ABUSE POTENTIAL (SISAP) QUESTIONNAIRE

1. If you drink, how many drinks do you have on a typical day?

If less than 5 for men/less than 4 for women, then ask question 2.

If 5 or more for men/4 or more for women, then you may stop here *Use caution when prescribing opioids.*

2. How many drinks do you have in a typical week?

If less than 17 for men/less than 13 for women, then ask question 3.

If 17 or more for men/13 or more for women, then you may stop here *Use caution when prescribing opioids.*

3. Have you used marijuana or hashish in the last year?

If no, then ask question 4.

If yes, then you may stop here *Use caution when prescribing opioids.*

4. Have you ever smoked cigarettes?

If no, then you may stop here *Probably a low opioid abuse risk.*

If yes, then ask question 5.

5. What is your age?

If under 40 years of age, then you may stop here *Use caution when prescribing opioids.*

If 40 years of age or older, then you may stop here *Probably a low opioid abuse risk.*

NOTES

- Use caution when prescribing opioids to these patients:
 - a. Men who drink more than 4 alcoholic beverages per day or 16 per week
 - b. Women who drink more than 3 alcoholic beverages per day or 12 per week
 - c. Persons who admit to recreational use of marijuana or hashish in the previous year
 - d. Persons who are younger than 40 years of age and smoke